

# Partial fractions II — repeated and quadratic factors

Lesson 73 · Module 3 — Integral Calculus · The Calculus Course · [dailymaths.uk/course](http://dailymaths.uk/course)

---

## Practice problems

1. Evaluate the integral  $\int \frac{1}{x^2-4x+4} dx$ .
2. Evaluate the integral  $\int \frac{1}{x^3+x} dx$ .
3. Evaluate the integral  $\int \frac{x^2}{x^2+1} dx$ .

## Solutions

1. Evaluate the integral  $\int \frac{1}{x^2-4x+4} dx$ .

- Factor the denominator:  $x^2 - 4x + 4 = (x - 2)^2$ .
- Set up partial fractions:  $\frac{1}{(x-2)^2} = \frac{A}{x-2} + \frac{B}{(x-2)^2}$ .
- Solve for constants:  $1 = A(x - 2) + B$ . Setting  $x = 2$  gives  $B = 1$ . Comparing coefficients of  $x$  gives  $A = 0$ .
- Integrate:  $\int \frac{1}{(x-2)^2} dx = -(x - 2)^{-1} + C$ .

**Answer:**  $-\frac{1}{x-2} + C$

2. Evaluate the integral  $\int \frac{1}{x^3+x} dx$ .

- Factor the denominator:  $x(x^2 + 1)$ .
- Set up partial fractions:  $\frac{1}{x(x^2+1)} = \frac{A}{x} + \frac{Bx+C}{x^2+1}$ .
- Solve for constants:  $1 = A(x^2 + 1) + (Bx + C)x$ .
- Comparing coefficients:  $x^2$  terms:  $A + B = 0$ ;  $x$  terms:  $C = 0$ ; constants:  $A = 1$ . Thus  $B = -1$ .
- Integrate:  $\int (\frac{1}{x} - \frac{x}{x^2+1}) dx = \ln|x| - \frac{1}{2} \ln(x^2 + 1) + C$ .

**Answer:**  $\ln|x| - \frac{1}{2} \ln(x^2+1) + C$

3. Evaluate the integral  $\int \frac{x^2}{x^2+1} dx$ .

- Check degree: Numerator degree is 2, denominator degree is 2. The fraction is improper.
- Perform long division:  $\frac{x^2}{x^2+1} = 1 - \frac{1}{x^2+1}$ .
- Integrate the resulting terms:  $\int 1 dx - \int \frac{1}{x^2+1} dx$ .
- Apply basic integration rules:  $x - \arctan(x) + C$ .

**Answer:**  $x - \arctan(x) + C$