

# Centre of mass and centroids

Lesson 92 · Module 4 — Applications of Integration · The Calculus Course · [dailymaths.uk/course](https://dailymaths.uk/course)

---

## Practice problems

1. Find the centroid of the region bounded by  $y = \sqrt{x}$ , the  $x$ -axis, and the line  $x = 4$ .
2. Find the centroid of the region bounded by  $y = x^2$  and  $y = x$ .
3. Find the centroid of the region bounded by  $y = e^x$ ,  $y = 0$ ,  $x = 0$ , and  $x = 1$ .

## Solutions

1. Find the centroid of the region bounded by  $y = \sqrt{x}$ , the  $x$ -axis, and the line  $x = 4$ .

- Calculate area:  $A = \int_0^4 x^{1/2} dx = [\frac{2}{3}x^{3/2}]_0^4 = \frac{2}{3}(8) = \frac{16}{3}$ .
- Calculate  $M_y$ :  $M_y = \int_0^4 x(x^{1/2}) dx = \int_0^4 x^{3/2} dx = [\frac{2}{5}x^{5/2}]_0^4 = \frac{2}{5}(32) = \frac{64}{5}$ .
- Calculate  $M_x$ :  $M_x = \int_0^4 \frac{1}{2}(\sqrt{x})^2 dx = \int_0^4 \frac{1}{2}x dx = [\frac{1}{4}x^2]_0^4 = 4$ .
- Find  $\bar{x} = \frac{64/5}{16/3} = \frac{64}{5} \cdot \frac{3}{16} = \frac{12}{5} = 2.4$ .
- Find  $\bar{y} = \frac{4}{16/3} = 4 \cdot \frac{3}{16} = \frac{3}{4} = 0.75$ .

**Answer:** ( $\frac{12}{5}$ ,  $\frac{3}{4}$ )

2. Find the centroid of the region bounded by  $y = x^2$  and  $y = x$ .

- Intersection points are  $x = 0$  and  $x = 1$ .
- Area  $A = \int_0^1 (x - x^2) dx = [\frac{1}{2}x^2 - \frac{1}{3}x^3]_0^1 = \frac{1}{6}$ .
- Moment  $M_y = \int_0^1 x(x - x^2) dx = [\frac{1}{3}x^3 - \frac{1}{4}x^4]_0^1 = \frac{1}{12}$ .
- Moment  $M_x = \int_0^1 \frac{1}{2}(x^2 - (x^2)^2) dx = \frac{1}{2}[\frac{1}{3}x^3 - \frac{1}{5}x^5]_0^1 = \frac{1}{2}(\frac{2}{15}) = \frac{1}{15}$ .
- $\bar{x} = \frac{1/12}{1/6} = \frac{1}{2}$ ,  $\bar{y} = \frac{1/15}{1/6} = \frac{2}{5}$ .

**Answer:** ( $\frac{1}{2}$ ,  $\frac{2}{5}$ )

3. Find the centroid of the region bounded by  $y = e^x$ ,  $y = 0$ ,  $x = 0$ , and  $x = 1$ .

- Area  $A = \int_0^1 e^x dx = [e^x]_0^1 = e - 1$ .
- Moment  $M_y = \int_0^1 xe^x dx$ . Using integration by parts:  $[xe^x - e^x]_0^1 = (e - e) - (0 - 1) = 1$ .
- Moment  $M_x = \int_0^1 \frac{1}{2}(e^x)^2 dx = \int_0^1 \frac{1}{2}e^{2x} dx = [\frac{1}{4}e^{2x}]_0^1 = \frac{1}{4}(e^2 - 1)$ .
- $\bar{x} = \frac{1}{e-1}$ .
- $\bar{y} = \frac{(e^2-1)/4}{e-1} = \frac{(e-1)(e+1)}{4(e-1)} = \frac{e+1}{4}$ .

**Answer:** ( $\frac{1}{e-1}$ ,  $\frac{e+1}{4}$ )