

Monotone functions and their discontinuities; inverse functions

Lesson 65 · Module 5 — Limits & Continuity · Real Analysis · dailymaths.uk/analysis

Practice problems

1. Let $f : [0, 1] \rightarrow \mathbb{R}$ be a monotone increasing function. Prove that the set $S = \{x \in (0, 1) : f(x^+) > f(x^-)\}$ is at most countable.
2. Let $f : [a, b] \rightarrow [c, d]$ be a strictly increasing continuous bijection. Prove that f^{-1} is continuous at any point $y_0 \in (c, d)$.
3. Construct a continuous bijection $f : D \rightarrow [0, 1]$ where D is not an interval, such that f^{-1} is not continuous.

Solutions

1. Let $f : [0, 1] \rightarrow \mathbb{R}$ be a monotone increasing function. Prove that the set $S = \{x \in (0, 1) : f(x^+) > f(x^-)\}$ is at most countable.
 - For each $x \in S$, define the jump interval $J_x = (f(x^-), f(x^+))$.
 - Since f is monotone increasing, if $x < y$, then $f(x^+) \leq f(y^-)$, which implies $J_x \cap J_y = \emptyset$.
 - Each J_x is a non-empty open interval, so it contains at least one rational number q_x .
 - The mapping $x \mapsto q_x$ is an injection from S into $\mathbb{Q}$.
 - Since $\mathbb{Q}$ is countable, S must be at most countable.

Answer: The set S is countable because it can be injected into the set of rational numbers.

2. Let $f : [a, b] \rightarrow [c, d]$ be a strictly increasing continuous bijection. Prove that f^{-1} is continuous at any point $y_0 \in (c, d)$.
 - Let $x_0 = f^{-1}(y_0)$. We want to show that for any $\epsilon > 0$, there exists $\delta > 0$ such that $|y - y_0| < \delta \implies |f^{-1}(y) - x_0| < \epsilon$.
 - Consider the interval $(x_0 - \epsilon, x_0 + \epsilon) \cap [a, b]$. This is an open set in the subspace topology of $[a, b]$.
 - Since f is strictly increasing and continuous on an interval, it maps the open interval $(x_0 - \epsilon, x_0 + \epsilon)$ to an open interval $(f(x_0 - \epsilon), f(x_0 + \epsilon))$.
 - Since $y_0 = f(x_0)$ is inside this open interval, there exists a $\delta > 0$ such that $(y_0 - \delta, y_0 + \delta) \subset (f(x_0 - \epsilon), f(x_0 + \epsilon))$.
 - Thus, if $|y - y_0| < \delta$, then $f^{-1}(y)$ must lie within $(x_0 - \epsilon, x_0 + \epsilon)$, proving continuity.

Answer: The continuity of f^{-1} follows from the fact that f maps open intervals to open intervals.

3. Construct a continuous bijection $f : D \rightarrow [0, 1]$ where D is not an interval, such that f^{-1} is not continuous.
 - Let $D = [0, 1/2] \cup [1, 3/2]$.
 - Define $f(x) = x$ for $x \in [0, 1/2]$ and $f(x) = x - 1/2$ for $x \in [1, 3/2]$.
 - The range of the first part is $[0, 1/2]$ and the range of the second part is $[1/2, 1]$.
 - The function is a bijection from D to $[0, 1]$ and is continuous on its domain.
 - The inverse is $f^{-1}(y) = y$ for $y \in [0, 1/2]$ and $f^{-1}(y) = y + 1/2$ for $y \in [1/2, 1]$.
 - As $y \rightarrow 1/2$ from the left, $f^{-1}(y) \rightarrow 1/2$, but $f^{-1}(1/2) = 1$. The jump at $y = 1/2$ proves discontinuity.

Answer: A function defined on $D = [0, 1/2] \cup [1, 3/2]$ with $f(x) = x$ on the first part and $f(x) = x - 1/2$ on the second part works.