

Module 3 Review and Self-Test

Lesson 46 · Module 3 — Combinatorics, Deepened · Discrete Mathematics · dailymaths.uk/discrete

Practice problems

1. Calculate the number of ways to partition a set of 6 distinct elements into 2 non-empty identical subsets.
2. Find the number of ways to triangulate a convex polygon with 6 sides.
3. A necklace has 3 beads and 3 available colors. How many distinct necklaces can be made if only rotations are allowed?

Solutions

1. Calculate the number of ways to partition a set of 6 distinct elements into 2 non-empty identical subsets.

- We need to find the Stirling number of the second kind $S(6, 2)$.
- The formula for $S(n, 2)$ is $2^{n-1} - 1$.
- Plugging in $n = 6$, we get $2^{6-1} - 1 = 2^5 - 1$.
- This equals $32 - 1 = 31$.

Answer: 31

2. Find the number of ways to triangulate a convex polygon with 6 sides.

- The number of ways to triangulate a convex polygon with $n + 2$ sides is given by the n -th Catalan number C_n .
- Here, $n + 2 = 6$, so $n = 4$.
- We calculate $C_4 = \frac{1}{4+1} \binom{2 \times 4}{4} = \frac{1}{5} \binom{8}{4}$.
- $\binom{8}{4} = \frac{8 \times 7 \times 6 \times 5}{4 \times 3 \times 2 \times 1} = 70$.
- The result is $70/5 = 14$.

Answer: 14

3. A necklace has 3 beads and 3 available colors. How many distinct necklaces can be made if only rotations are allowed?

- The group G has 3 rotations: $0^\circ, 120^\circ, 240^\circ$.
- For 0° , all $3^3 = 27$ colorings are fixed.
- For 120° , all beads must be the same color, so 3 fixed points.
- For 240° , all beads must be the same color, so 3 fixed points.
- By Burnside's Lemma, the number of orbits is $\frac{1}{3}(27 + 3 + 3) = \frac{33}{3} = 11$.

Answer: 11